arXiv:1701.03725v1 [math.NT] 4 Jan 2017

Explicit Evaluations of Sums of Sequence Tails

Ce Xu* Xiaolian Zhou†

School of Mathematical Sciences, Xiamen University

Xiamen 361005, P.R. China

Abstract In this paper, we use Abel's summation formula to evaluate several quadratic sums of the form:

$$F_N(A,B;x) := \sum_{n=1}^{N} (A-A_n)(B-B_n)x^n,\ x\in[-1,1],$$

where the sequences A_n, B_n are defined by the sums

$$A_n = \sum_{k=1}^{n} a_k, B_n = \sum_{k=1}^{n} b_k,\quad a_k, b_k = o(n^{-p}), \Re(p) > 1$$

and $A = \lim_{n\to\infty} A_n, B = \lim_{n\to\infty} B_n, F(A,B;x) = \lim_{n\to\infty} F_n(A,B;x)$. We give an explicit formula of $F_n(A,B;x)$. Furthermore, we also evaluate several other series involving multiple zeta star values.

Keywords Sequence; harmonic numbers; Abel's summation formula; Riemann zeta function; multiple zeta star values, tails.

AMS Subject Classifications (2010): 11M06; 11M32.

1 Introduction

In recent years there has been significant interest in investigating sums of products of Riemann zeta tails and multiple zeta values. The subject of this paper is studying quadratic and cubic sums

$$F(A,B;x) = \sum_{n=1}^{\infty} (A-A_n)(B-B_n)x^n, F(A,B,\zeta(r)) = \sum_{n=1}^{\infty} (A-A_n)(B-B_n)(\zeta(r)-\zeta_n(r)),$$

where

$$H_n = \zeta_n(1) = \sum_{j=1}^{n} \frac{1}{j},\ \zeta_n(r) = \sum_{j=1}^{n} \frac{1}{j^r},\ r,n\in\mathbb{N} = \{1,2,3\cdots\}$$

are classical harmonic number and generalized harmonic numbers or partial sums of the series

$$\zeta(s) = \sum_{j=1}^{\infty} \frac{1}{j^s},\ \Re(s) > 1$$

*Corresponding author. Email: 15959259051@163.com

†Email: xlzhou@stu.xmu.edu.cn (X. ZHOU)

defining the Riemann zeta function, respectively, where the empty sum $\zeta_0(r)$ is conventionally understood to be zero. In recent papers [9,10], O. Furdui, A. Sîntămărian and C. Vălean prove some results on sums of products of the tails of the Riemann zeta function, i.e.,

$$\zeta(s) - \zeta_n(s) = \sum_{j=n+1}^{\infty} \frac{1}{j^s}.$$

In [9], O. Furdui and A. Sîntămărian shown that quadratic series

$$F(\zeta(s), \zeta(s)) = \sum_{n=1}^{\infty} (\zeta(s) - \zeta_n(s))^2, \ s \in \mathbb{N}/\{1\}$$

can be expressed as a rational linear combination of zeta function values, binomial coefficients and products of two zeta function values. In [10], O. Furdui and C. Vălean proven that the series

$$F(\zeta(s), \zeta(s+1)) = \sum_{n=1}^{\infty} (\zeta(s) - \zeta_n(s))(\zeta(s+1) - \zeta_n(s+1)), \ s \in \mathbb{N}/\{1\}$$

can be expressed in terms of Riemann zeta function values and a special integral involving a polylogarithm function. The polylogarithm function defined as follows

$$\mathrm{Li}_p(x) = \sum_{n=1}^{\infty} \frac{x^n}{n^p}, \Re(p) > 1, \ |x| \leq 1.$$

Furthermore, in [12], M. E. Hoffman shown a simple general result for the quadratic sum

$$F(\zeta(p), \zeta(q)) = \sum_{n=1}^{\infty} (\zeta(p) - \zeta_n(p))(\zeta(q) - \zeta_n(q)), \ p, q \in \mathbb{N}/\{1\}$$

in terms of the multiple zeta values $\zeta(s_1, s_2, \ldots, s_m)$ defined by

$$\zeta(s_1, s_2, \ldots, s_m) = \sum_{k_1 > \cdots > k_m \geq 1} \frac{1}{k_1^{s_1} \cdots k_m^{s_m}},$$

where for convergence $s_1 + s_2 + \cdots + s_j > j$ for $j = 1, 2, \ldots, m$. Moreover, Hoffman also provided an explicit evaluation of sums of products of three or more Riemann zeta tails in a closed form in terms of multiple zeta values (see Theorem 4.3 in the reference [12]). The multiple zeta value $\zeta(s_1, s_2, \ldots, s_m)$ is said to have depth m and weight $s_1 + \ldots + s_m$. Multiple zeta values of general depth were introduced independently by Hoffman [11] and D. Zagier [19], but for depth two they were already studied by Euler. In [5], Borwein and Girgensohn gave some evaluation multiple zeta values of depth three.

In this paper, we discuss the analytic representations of $F(A, B; x)$ and $F(A, B, \zeta(r))$. Using certain integral representations, we can prove that the cubic series

$$F(\zeta(p), \zeta(q), \zeta(r)) = \sum_{n=1}^{\infty} (\zeta(p) - \zeta_n(p))(\zeta(q) - \zeta_n(q))(\zeta(r) - \zeta_n(r)), \ p, q, r \in \mathbb{N}/\{1\}$$

can be expressed in terms of multiple zeta star values defined by

$$\zeta^{\star}(s_1, s_2, \ldots, s_m) = \sum_{k_1 \geq \cdots \geq k_m \geq 1} \frac{1}{k_1^{s_1} \cdots k_m^{s_m}}, \ s_1 \in \mathbb{N}/\{1\}, s_i \in \mathbb{N}, (i = 2, 3, \ldots).$$

Obviously, $\zeta^{\star}(m,p)=\sum\limits_{n=1}^{\infty}\dfrac{\zeta_n(p)}{n^m}$. Furthermore, we also give some evaluation of sums of products of alternating Riemann Zeta Tails. For example,

$$\begin{aligned}&\sum_{n=1}^{\infty}\left(\zeta(m)-\zeta_n(m)\right)\left(\bar{\zeta}(p)-\bar{\zeta}_n(p)\right)\\&=\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m-1)}{n^p}(-1)^{n-1}+\frac{\bar{\zeta}_n(p-1)}{n^m}\right\}-\zeta(m)\,\bar{\zeta}(p)-\bar{\zeta}(m+p-1),\end{aligned}$$

where $\bar{\zeta}(p)$ and $\bar{\zeta}_n(p)$ stand for the alternating Riemann Zeta function and alternating harmonic numbers defined by

$$\bar{\zeta}(p)=\sum_{k=1}^{\infty}\frac{(-1)^{k-1}}{k^p},\bar{\zeta}_n(p)=\sum_{k=1}^{n}\frac{(-1)^{k-1}}{k^p},n\in\mathbb{N},\Re(p)\geq 1.$$

The evaluation of linear sums$\sum\limits_{n=1}^{\infty}\dfrac{\zeta_n(p)}{n^m}$,$\sum\limits_{n=1}^{\infty}\dfrac{\bar{\zeta}_n(p)}{n^m}(-1)^{n-1}$,$\sum\limits_{n=1}^{\infty}\dfrac{\zeta_n(p)}{n^m}(-1)^{n-1}$ and $\sum\limits_{n=1}^{\infty}\dfrac{\bar{\zeta}_n(p)}{n^m}$ in terms of values of Riemann zeta function (or alternating Riemann Zeta function) and polylogarithm function at positive integers is known when $(p,m)=(1,3),(2,2)$ or $p+m$ is odd (see [2,3,7,14,17,18]). For example,

$$\begin{aligned}&\sum_{n=1}^{\infty}\frac{\bar{\zeta}_n(1)}{n^3}=\frac{7}{4}\zeta(3)\ln 2-\frac{5}{16}\zeta(4),\\&\sum_{n=1}^{\infty}\frac{H_n}{n^3}(-1)^{n-1}=-2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{11}{4}\zeta(4)+\frac{1}{2}\zeta(2)\ln^2 2-\frac{1}{12}\ln^4 2-\frac{7}{4}\zeta(3)\ln 2,\\&\sum_{n=1}^{\infty}\frac{\bar{\zeta}_n(1)}{n^3}(-1)^{n-1}=\frac{3}{2}\zeta(4)+\frac{1}{2}\zeta(2)\ln^2 2-\frac{1}{12}\ln^4 2-2\mathrm{Li}_4\left(\frac{1}{2}\right),\\&\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^2}(-1)^{n-1}=-\frac{51}{16}\zeta(4)+4\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{7}{2}\zeta(3)\ln 2-\zeta(2)\ln^2(2)+\frac{1}{6}\ln^4 2.\end{aligned}$$

Furthermore, we consider the nested sum

$$\sum_{n=1}^{\infty}\frac{y^n}{n^m}\left(\sum_{k=1}^{n}\frac{x^k}{k^p}\right),\ x,y\in[-1,1),\ m,p\in\mathbb{N}.$$

By taking the sum over complementary pairs of summation indices, we obtain a simple reflection formula

$$\sum_{n=1}^{\infty}\frac{y^n}{n^m}\left(\sum_{k=1}^{n}\frac{x^k}{k^p}\right)+\sum_{n=1}^{\infty}\frac{x^n}{n^p}\left(\sum_{k=1}^{n}\frac{y^k}{k^m}\right)=\mathrm{Li}_p(x)\,\mathrm{Li}_m(y)+\mathrm{Li}_{p+m}(xy).$$

Therefore, we can deduce that

$$\sum_{n=1}^{\infty}\frac{\bar{\zeta}_n(2)}{n^2}(-1)^{n-1}=\frac{13}{16}\zeta(4),$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n(3)}{n}(-1)^{n-1}=\frac{19}{16}\zeta(4)-\frac{3}{4}\zeta(3)\ln 2,$$
$$\sum_{n=1}^{\infty}\frac{\bar{\zeta}_n(2)}{n^2}=\frac{85}{16}\zeta(4)-4\mathrm{Li}_4\left(\frac{1}{2}\right)+\zeta(2)\ln^2 2-\frac{1}{6}\ln^4 2-\frac{7}{2}\zeta(3)\ln 2,$$
$$\sum_{n=1}^{\infty}\frac{\bar{\zeta}_n(3)}{n}(-1)^{n-1}=2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{3}{4}\zeta(3)\ln 2+\frac{1}{12}\ln^4 2-\frac{1}{2}\zeta(4)-\frac{1}{2}\zeta(2)\ln^2 2.$$

Some results for sums of harmonic numbers or alternating harmonic numbers may be seen in the works of [2,3,7,13-18] and references therein.
The main result of this paper is the following theorems.

Theorem 1.1 *If $a_k, b_k = o(n^{-p}), \Re(p) > 1$ and $x \in [-1,1]$, then*

$$\begin{aligned}(1-x)F_n(A,B;x)&=(1-x)\sum_{k=1}^{n}(A-A_k)(B-B_k)x^k\\&=ABx-(A-A_n)(B-B_n)x^{n+1}-\left(\sum_{k=1}^{n}a_kx^k\right)(B-B_n)-\left(\sum_{k=1}^{n}b_kx^k\right)(A-A_n)\\&\quad-\sum_{k=1}^{n}\left(\sum_{i=1}^{k}a_ix^i\right)b_k-\sum_{k=1}^{n}\left(\sum_{i=1}^{k}b_ix^i\right)a_k+\sum_{k=1}^{n}a_kb_kx^k.\end{aligned}\tag{1.1}$$

Theorem 1.2 *If $a_k, b_k = o(n^{-p}), \Re(p) > 1$ and $r \in \mathbb{N}/\{1\}$, then*

$$\begin{aligned}F(A,B,\zeta(r))=&\sum_{k=1}^{\infty}(A-A_k)(B-B_k)(\zeta(r)-\zeta_k(r))\\=&\zeta(r)\left\{\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}ka_k\right)b_n+\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}kb_k\right)a_n-\sum_{n=1}^{\infty}na_nb_n-AB\right\}\\&-\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}a_k(k\zeta_k(r)-\zeta_k(r-1))\right)b_n-\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}b_k(k\zeta_k(r)-\zeta_k(r-1))\right)a_n\\&+\sum_{n=1}^{\infty}a_nb_n(n\zeta_n(r)-\zeta_n(r-1)).\end{aligned}\tag{1.2}$$

In section 2, we will use Abels summation formula to prove Theorem 1.1 and 1.2. The Abels summation by parts formula states that if $(a_n)_{n\geq 1}$ and $(b_n)_{n\geq 1}$ are two sequences of real numbers and $A_n=\sum_{k=1}^{n}a_k$, then $\sum_{k=1}^{n}a_kb_k=A_nb_{n+1}+\sum_{k=1}^{n}A_k(b_k-b_{k+1})$.

2 Proof of Theorem 1.1 and Theorem 1.2

We now prove our Theorems. Firstly, we are ready to prove Theorem 1.1.
Proof. From the definition $F_n(A,B;x)$, we have

$$F_n(A,B;x)=\sum_{k=1}^{n}(A-A_k)(B-B_k)x^k$$

$$
\begin{aligned}
&= (A-a_1)(B-b_1)x + \sum_{k=1}^{n-1} (A-A_{k+1})(B-B_{k+1}) x^{k+1} \\
&= (A-a_1)(B-b_1)x + \sum_{k=1}^{n-1} (A-A_k-a_{k+1})(B-B_k-b_{k+1}) x^{k+1} \\
&= (A-a_1)(B-b_1)x + \sum_{k=1}^{n-1} (A-A_k)(B-B_k) x^{k+1} \\
&\quad - \sum_{k=1}^{n-1} \{a_{k+1}(B-B_k) + b_{k+1}(A-A_k)\} x^{k+1} + \sum_{k=1}^{n-1} a_{k+1}b_{k+1}x^{k+1} \\
&= (A-a_1)(B-b_1)x + \sum_{k=1}^{n} (A-A_k)(B-B_k) x^{k+1} - (A-A_n)(B-B_n)x^{n+1} \\
&\quad + \sum_{k=1}^{n} a_k b_k x^k - a_1 b_1 x - \sum_{k=1}^{n} \{a_k(B-B_k+b_k) + b_k(A-A_k+a_k)\} x^k + a_1 B + b_1 A \\
&= xF_n(A,B;x) + ABx - (A-A_n)(B-B_n)x^{n+1} \\
&\quad - \sum_{k=1}^{n} \{a_k(B-B_k) + b_k(A-A_k)\} x^k - \sum_{k=1}^{n} a_k b_k x^k. \qquad (2.1)
\end{aligned}
$$

By Abel's summation formula, we can deduce that

$$
\sum_{k=1}^{n} \{a_k(B-B_k)\} x^k = \left(\sum_{k=1}^{n} a_k x^k\right)(B-B_n) + \sum_{k=1}^{n} \left(\sum_{i=1}^{k} a_i x^i\right) b_k - \sum_{k=1}^{n} a_k b_k x^k, \qquad (2.2)
$$

$$
\sum_{k=1}^{n} \{b_k(A-A_k)\} x^k = \left(\sum_{k=1}^{n} b_k x^k\right)(A-A_n) + \sum_{k=1}^{n} \left(\sum_{i=1}^{k} b_i x^i\right) a_k - \sum_{k=1}^{n} a_k b_k x^k. \qquad (2.3)
$$

Substituting (2.2) (2.3) into (2.1) respectively, we can obtain (1.1). We have completed the proof of Theorem 1.1. Now, we are ready to prove Theorem 1.2. Multiplying (1.2) by $\dfrac{\ln^{r-1}x}{(1-x)^2}$ and integrating over the interval (0,1), and using the following elementary integral identities

$$
\int_0^1 \frac{x^k \ln^{r-1}x}{1-x} dx = (-1)^{r-1}(r-1)!\left(\zeta(r) - \zeta_k(r)\right),\ r \in \mathbb{N}/\{1\},
$$

$$
\int_0^1 \frac{x^k \ln^{r-1}x}{(1-x)^2} dx = (-1)^{r-1}(r-1)!\left(\zeta(r-1) - \zeta_k(r-1) - k\zeta(r) + k\zeta_k(r)\right), r \in \mathbb{N}/\{1,2\},
$$

by a direct calculation, we obtain

$$
\begin{aligned}
F(A,B,\zeta(r)) =& AB(\zeta(r-1) - \zeta(r)) \\
&+ \sum_{k=1}^{\infty} a_k b_k (\zeta(r-1) - \zeta_k(r-1) - k\zeta(r) + k\zeta_k(r))
\end{aligned}
$$

$$-\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}a_k\left(\zeta\left(r-1\right)-\zeta_k\left(r-1\right)-k\zeta\left(r\right)+k\zeta_k\left(r\right)\right)\right)b_n$$
$$-\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}b_k\left(\zeta\left(r-1\right)-\zeta_k\left(r-1\right)-k\zeta\left(r\right)+k\zeta_k\left(r\right)\right)\right)a_n. \tag{2.4}$$

Taking $x=1$ in (1.1), we get

$$\sum_{k=1}^{n}A_kb_k+\sum_{k=1}^{n}B_ka_k=\sum_{k=1}^{n}a_kb_k+A_nB_n. \tag{2.5}$$

Letting $n\to\infty$, then have

$$\sum_{k=1}^{\infty}A_kb_k+\sum_{k=1}^{\infty}B_ka_k=\sum_{k=1}^{\infty}a_kb_k+AB. \tag{2.6}$$

Substituting (2.6) into (2.4), we know that when $r\in\mathbb{N}/\{1,2\}$, the equality (1.2) is holds. Similarly, multiplying (1.2) by $\dfrac{\ln x}{(1-x)^2}$ and integrating over the interval (0,t), then letting $t\to 1$, we can prove that if $r=2$, the formula (1.2) is also true. The proof of Theorem 1.2 is finished.

3 Some results

In this section, we will give some closed form of sums of sequences tails.
Firstly, multiplying $(1-x)^{-1}$ in both sides of (1.1), then letting $x\to 1$ and using L. Hopital's rule, we deduce that

$$\sum_{k=1}^{n}\left(A-A_k\right)\left(B-B_k\right)=F_n\left(A,B;1\right)=\lim_{x\to 1}\frac{(1-x)F_n\left(A,B;x\right)}{1-x}$$
$$=\sum_{k=1}^{n}\left(\sum_{i=1}^{k}ia_i\right)b_k+\sum_{k=1}^{n}\left(\sum_{i=1}^{k}ib_i\right)a_k+\left(\sum_{k=1}^{n}ka_k\right)\left(B-B_n\right)+\left(\sum_{k=1}^{n}kb_k\right)\left(A-A_n\right)$$
$$+\left(n+1\right)\left(A-A_n\right)\left(B-B_n\right)-\left(\sum_{k=1}^{n}ka_kb_k\right)-AB. \tag{3.1}$$

Letting $n\to\infty$ in (3.1), we have

$$\sum_{n=1}^{\infty}\left(A-A_n\right)\left(B-B_n\right)=\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}ka_k\right)b_n+\sum_{n=1}^{\infty}\left(\sum_{k=1}^{n}kb_k\right)a_n-\sum_{n=1}^{\infty}na_nb_n-AB. \tag{3.2}$$

Putting $(a_n,b_n)=\left(\dfrac{1}{n^m},\dfrac{1}{n^p}\right),\left(\dfrac{(-1)^{n-1}}{n^m},\dfrac{(-1)^{n-1}}{n^p}\right),\left(\dfrac{1}{n^m},\dfrac{(-1)^{n-1}}{n^p}\right)$ in (3.2), we can obtain the following identities

$$\sum_{n=1}^{\infty}\left(\zeta\left(m\right)-\zeta_n\left(m\right)\right)\left(\zeta\left(p\right)-\zeta_n\left(p\right)\right)$$

$$= \zeta^{\star}(p, m-1) + \zeta^{\star}(m, p-1) - \zeta(m+p-1) - \zeta(m)\zeta(p), \tag{3.3}$$

$$\sum_{n=1}^{\infty} \left(\bar{\zeta}(m) - \bar{\zeta}_n(m)\right)\left(\bar{\zeta}(p) - \bar{\zeta}_n(p)\right)$$

$$= \sum_{n=1}^{\infty} \frac{\bar{\zeta}_n(p-1)}{n^m}(-1)^{n-1} + \sum_{n=1}^{\infty} \frac{\bar{\zeta}_n(m-1)}{n^p}(-1)^{n-1} - \zeta(m+p-1) - \bar{\zeta}(m)\bar{\zeta}(p), \tag{3.4}$$

$$\sum_{n=1}^{\infty} \left(\zeta(m) - \zeta_n(m)\right)\left(\bar{\zeta}(p) - \bar{\zeta}_n(p)\right)$$

$$= \sum_{n=1}^{\infty} \left\{ \frac{\zeta_n(m-1)}{n^p}(-1)^{n-1} + \frac{\bar{\zeta}_n(p-1)}{n^m} \right\} - \zeta(m)\bar{\zeta}(p) - \bar{\zeta}(m+p-1). \tag{3.5}$$

Similarly, taking $(a_n, b_n) = \left(\frac{1}{n^m}, \frac{1}{n^p}\right)$ in (1.2), we arrive at the conclusion that

$$\begin{aligned} F(\zeta(m), \zeta(p), \zeta(r)) =& \sum_{n=1}^{\infty} \left(\zeta(m) - \zeta_n(m)\right)\left(\zeta(p) - \zeta_n(p)\right)\left(\zeta(r) - \zeta_n(r)\right) \\ =& \zeta(r)\left\{\zeta^{\star}(p, m-1) + \zeta^{\star}(m, p-1) - \zeta(m+p-1) - \zeta(m)\zeta(p)\right\} \\ &+ \left\{\zeta^{\star}(m+p-1, r) - \zeta^{\star}(m+p, r-1)\right\} \\ &- \left\{\zeta^{\star}(p, m-1, r) - \zeta^{\star}(p, m, r-1)\right\} \\ &- \left\{\zeta^{\star}(m, p-1, r) - \zeta^{\star}(m, p, r-1)\right\}. \end{aligned} \tag{3.6}$$

In (1.1), letting $n \to \infty$, then multiplying it by $\dfrac{\ln^{r-1} x}{x(1-x)}$ and integrating over the interval (0,1), we conclude that

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{(A - A_n)(B - B_n)}{n^r} =& \sum_{n=1}^{\infty} \left(\sum_{k=1}^{n} a_k \zeta_{k-1}(r)\right) b_n + \sum_{n=1}^{\infty} \left(\sum_{k=1}^{n} b_k \zeta_{k-1}(r)\right) a_n \\ &- \sum_{n=1}^{\infty} a_n b_n \zeta_{n-1}(r), \qquad\qquad r \in \mathbb{N}/\{1\}. \end{aligned} \tag{3.7}$$

In fact, it is easily shown that the formula (3.7) is also true when $r = 1$.
Taking $(a_n, b_n) = \left(\frac{1}{n^m}, \frac{1}{n^p}\right)$ in (3.7), we get

$$\begin{aligned} &\sum_{n=1}^{\infty} \frac{\left(\zeta(m) - \zeta_n(m)\right)\left(\zeta(p) - \zeta_n(p)\right)}{n^r} \\ &= \zeta(m+p+r) + \zeta^{\star}(m, p, r) + \zeta^{\star}(p, m, r) \\ &\quad - \zeta^{\star}(m+p, r) - \zeta^{\star}(m, p+r) - \zeta^{\star}(p, m+r), m, p, r \in \mathbb{N}. \end{aligned} \tag{3.8}$$

We define the multiple harmonic star number $\zeta_n^{\star}(s_1, s_2, \ldots, s_m)$ by

$$\zeta_n^{\star}(s_1, s_2, \ldots, s_m) := \sum_{1 \le k_m \le \cdots \le k_1 \le n} \frac{1}{k_1^{s_1} \cdots k_m^{s_m}}, s_i \in \mathbb{N},\ (i = 1, 2, \ldots, m).$$

From the definition multiple harmonic star number, we can deduce the following relation

$$\zeta^{\star}(p,m,r)+\sum_{n=1}^{\infty}\frac{\zeta_n(p)\,\zeta_n(r)}{n^m}=\zeta^{\star}(p+m,r)+\zeta(p)\,\zeta^{\star}(m,r),\quad p,m\in\mathbb{N}/\{1\},r\in\mathbb{N}. \tag{3.9}$$

Substituting (3.9) into (3.8) respectively, we obtain the symmetry identity

$$\begin{aligned}&\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m)\,\zeta_n(p)}{n^r}+\frac{\zeta_n(p)\,\zeta_n(r)}{n^m}+\frac{\zeta_n(m)\,\zeta_n(r)}{n^p}\right\}\\&=\zeta^{\star}(p+m,r)+\zeta^{\star}(p+r,m)+\zeta^{\star}(m+r,p)\\&\quad+\zeta(p)\,\zeta(m)\,\zeta(r)-\zeta(p+m+r),\qquad p,m,r\in\mathbb{N}/\{1\}.\end{aligned} \tag{3.10}$$

From [7,17], we know that the quadratic sums $\sum_{n=1}^{\infty}\frac{\zeta_n(m)\,\zeta_n(p)}{n^r}$ can be evaluated in terms of zeta values and double zeta star values $\zeta^{\star}(s_1,s_2)$ in the following cases: $m=p=r$ or $m+p+r$ even $(m,p,r\in\mathbb{N}/\{1\})$ and $m=1,p+r\leq 8$. From example

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_n\zeta_n(3)}{n^3}&=\frac{83}{8}\zeta(7)+\frac{1}{4}\zeta(3)\,\zeta(4)-\frac{11}{2}\zeta(2)\,\zeta(5),\\\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n^5}&=-\frac{343}{48}\zeta(8)+12\zeta(3)\,\zeta(5)-\frac{5}{2}\zeta(2)\,\zeta^2(3)-\frac{3}{4}\zeta^{\star}(6,2),\\\sum_{n=1}^{\infty}\frac{\zeta_n^2(2)}{n^4}&=11\zeta^{\star}(6,2)+\frac{457}{18}\zeta(8)+6\zeta(2)\,\zeta^2(3)-40\zeta(3)\,\zeta(5).\end{aligned}$$

where $\zeta^{\star}(6,2)=\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^6}$.

Similarly, from (1.1), we have the following results

$$\begin{aligned}&\sum_{n=1}^{\infty}\left(\zeta(m)-\zeta_n(m)\right)\left(\zeta(p)-\zeta_n(p)\right)(-1)^{n-1}\\&=\frac{1}{2}\left\{\zeta(m)\,\zeta(p)+\bar{\zeta}(m+p)-\sum_{n=1}^{\infty}\frac{\bar{\zeta}_n(p)}{n^m}-\sum_{n=1}^{\infty}\frac{\bar{\zeta}_n(m)}{n^p}\right\},\end{aligned} \tag{3.11}$$

$$\begin{aligned}&\sum_{n=1}^{\infty}\left(\zeta^{\star}(p+1,p)-\zeta_n^{\star}(p+1,p)\right)\left(\zeta^{\star}(q+1,q)-\zeta_n^{\star}(q+1,q)\right)\\&=\frac{1}{2}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n^2(p)\,\zeta_n(q)+\zeta_n(2p)\,\zeta_n(q)}{n^{q+1}}+\frac{\zeta_n^2(q)\,\zeta_n(p)+\zeta_n(2q)\,\zeta_n(p)}{n^{p+1}}\right\}\\&\quad-\sum_{n=1}^{\infty}\frac{\zeta_n(p)\,\zeta_n(q)}{n^{p+q+1}}-\left(\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^{p+1}}\right)\left(\sum_{n=1}^{\infty}\frac{\zeta_n(q)}{n^{q+1}}\right),\end{aligned} \tag{3.12}$$

$$\begin{aligned}&\sum_{n=1}^{\infty}\left(\zeta^{\star}(p+1,p)-\zeta_n^{\star}(p+1,p)\right)\left(\zeta^{\star}(p+2,p)-\zeta_n^{\star}(p+2,p)\right)\\&=\frac{1}{2}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n^3(p)+\zeta_n(p)\,\zeta_n(2p)}{n^{p+2}}-\frac{\zeta_n^2(p)}{n^{2p+2}}\right\}+\frac{1}{2}\left(\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^{p+1}}\right)^2\end{aligned}$$

$$-\left(\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^{p+1}}\right)\left(\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^{p+2}}\right). \tag{3.13}$$

Next, we give a Theorem.

Theorem 3.1 *Let $m,p\geq 2$ be integers, then have*

$$\sum_{n=1}^{\infty}\frac{\zeta(m)\zeta_n(p)-\zeta(p)\zeta_n(m)}{n}=\zeta(p)\sum_{n=1}^{\infty}\frac{H_n}{n^m}-\zeta(m)\sum_{n=1}^{\infty}\frac{H_n}{n^p}+\zeta(m)\zeta(p+1)-\zeta(p)\zeta(m+1). \tag{3.14}$$

Proof. First, we construct the following power function

$$y=\sum_{n=1}^{\infty}\left\{H_n\zeta_n(m)-\zeta_n(m+1)\right\}x^{n-1},\ x\in(-1,1).$$

By the definition of harmonic numbers, we can deduce that

$$\begin{aligned}y&=\sum_{n=1}^{\infty}\left\{\left(H_n+\frac{1}{n+1}\right)\left(\zeta_n(m)+\frac{1}{(n+1)^m}\right)-\left(\zeta_n(m+1)+\frac{1}{(n+1)^{m+1}}\right)\right\}x^n\\&=\sum_{n=1}^{\infty}\left\{H_n\zeta_n(m)-\zeta_n(m+1)+\frac{H_n}{(n+1)^m}+\frac{\zeta_n(m)}{n+1}\right\}x^n.\end{aligned} \tag{3.15}$$

Therefore, from (3.15), we obtain the following relation

$$\sum_{n=1}^{\infty}\left\{H_n\zeta_n(m)-\zeta_n(m+1)\right\}x^{n-1}=\sum_{n=1}^{\infty}\left\{\frac{H_n}{(n+1)^m}+\frac{\zeta_n(m)}{n+1}\right\}\frac{x^n}{1-x}. \tag{3.16}$$

Then multiplying (3.16) by $\ln^{p-1}x$ and integrating over (0,1), we have

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(m)-\zeta_n(m+1)}{n^p}=\sum_{n=1}^{\infty}\left\{\frac{H_n}{(n+1)^m}+\frac{\zeta_n(m)}{n+1}\right\}\{\zeta(p)-\zeta_n(p)\}. \tag{3.17}$$

By a direct calculation, we obtain that

$$\begin{aligned}\sum_{n=1}^{\infty}\left\{\frac{H_n\zeta_n(m)}{n^p}+\frac{H_n\zeta_n(p)}{n^m}\right\}&=\zeta(p)\sum_{n=1}^{\infty}\frac{H_n}{n^m}+\sum_{n=1}^{\infty}\frac{H_n}{n^{p+m}}+\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n^{p+1}}\\&\quad-\sum_{n=1}^{\infty}\frac{\zeta_n(m+1)}{n^p}+\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n}\{\zeta(p)-\zeta_n(p)\}.\end{aligned} \tag{3.18}$$

Change m,p to p,m, we get

$$\begin{aligned}\sum_{n=1}^{\infty}\left\{\frac{H_n\zeta_n(p)}{n^m}+\frac{H_n\zeta_n(m)}{n^p}\right\}&=\zeta(m)\sum_{n=1}^{\infty}\frac{H_n}{n^p}+\sum_{n=1}^{\infty}\frac{H_n}{n^{m+p}}+\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^{m+1}}\\&\quad-\sum_{n=1}^{\infty}\frac{\zeta_n(p+1)}{n^m}+\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n}\{\zeta(m)-\zeta_n(m)\}.\end{aligned} \tag{3.19}$$

Combining (3.18) and (3.19), we can obtain (3.14). □

We note that

$$\sum_{n=1}^{\infty}\frac{\zeta(m)\zeta(p)-\zeta_n(m)\zeta_n(p)}{n}$$
$$=\zeta(p)\sum_{n=1}^{\infty}\frac{\zeta(m)-\zeta_n(m)}{n}+\sum_{n=1}^{\infty}\frac{\zeta_n(m)(\zeta(p)-\zeta_n(p))}{n},\ m,p\in\mathbb{N}/\{1\}. \tag{3.20}$$

In [10], O. Furdui and C. Valean gave the following formula

$$\sum_{n=1}^{\infty}\frac{\zeta(m)-\zeta_n(m)}{n}=\zeta^{\star}(m,1)-\zeta(m+1). \tag{3.21}$$

Hence, combining (3.18), (3.20) and (3.21), we have the following corollary.

Corollary 3.2 *For positive integers $m>1,p>1$, then we have*

$$\sum_{n=1}^{\infty}\frac{\zeta(m)\zeta(p)-\zeta_n(m)\zeta_n(p)}{n}$$
$$=\sum_{n=1}^{\infty}\left\{\frac{H_n\zeta_n(m)}{n^p}+\frac{H_n\zeta_n(p)}{n^m}\right\}+\zeta^{\star}(p,m+1)$$
$$-\zeta(p)\zeta(m+1)-\zeta^{\star}(p+m,1)-\zeta^{\star}(p+1,m). \tag{3.22}$$

Similarly, noting that

$$\sum_{n=1}^{\infty}\frac{\zeta_n(m)\zeta(p)\zeta(q)-\zeta(m)\zeta_n(p)\zeta_n(q)}{n}$$
$$=\zeta(m)\sum_{n=1}^{\infty}\frac{\zeta(m)\zeta(p)-\zeta_n(m)\zeta_n(p)}{n}-\zeta(p)\zeta(q)\sum_{n=1}^{\infty}\frac{\zeta(m)-\zeta_n(m)}{n},\quad m,p,q\in\mathbb{N}/\{1\}.$$

Therefore, we also obtain the following corollary.

Corollary 3.3 *For positive integers $m,p,q>1$, then we have*

$$\sum_{n=1}^{\infty}\frac{\zeta_n(m)\zeta(p)\zeta(q)-\zeta(m)\zeta_n(p)\zeta_n(q)}{n}$$
$$=\zeta(m)\left\{\begin{array}{l}\displaystyle\sum_{n=1}^{\infty}\left(\frac{H_n\zeta_n(q)}{n^p}+\frac{H_n\zeta_n(p)}{n^q}\right)+\zeta^{\star}(p,q+1)\\ -\zeta^{\star}(p+q,1)-\zeta^{\star}(p+1,q)-\zeta(p)\zeta(q+1)\end{array}\right\}$$
$$-\zeta(p)\zeta(q)\{\zeta^{\star}(m,1)-\zeta(m+1)\}. \tag{3.23}$$

Furthermore, we consider the following power series function

$$y=\sum_{n=1}^{\infty}\{H_n\zeta_n(p)\zeta_n(q)-\zeta_n(p+q+1)\}x^{n-1},\ x\in(-1,1),$$

proceeding in a similar fashion to evaluation of the Theorem 3.1, we can give the following Theorem.

Theorem 3.4 *For positive integers $p, m > 1$ and $q > 0$, we have*

$$\begin{aligned}
&\sum_{n=1}^{\infty} \frac{\zeta(m)\,\zeta_n(p)\,\zeta_n(q) - \zeta(p)\,\zeta_n(m)\,\zeta_n(q)}{n} \\
&= \zeta(p)\left\{\sum_{n=1}^{\infty}\left(\frac{H_n\zeta_n(m)}{n^q} + \frac{H_n\zeta_n(q)}{n^m}\right) - \zeta^\star(q+1,m) - \zeta^\star(m+q,1) - \zeta^\star(m+1,q)\right\} \\
&\quad - \zeta(m)\left\{\sum_{n=1}^{\infty}\left(\frac{H_n\zeta_n(q)}{n^p} + \frac{H_n\zeta_n(p)}{n^q}\right) - \zeta^\star(q+1,p) - \zeta^\star(p+q,1) - \zeta^\star(p+1,q)\right\} \\
&\quad + \zeta(m+q+1)\,\zeta(p) - \zeta(p+q+1)\,\zeta(m).
\end{aligned}$$

4 Some examples

Some illustrative examples follow.

$$\sum_{n=1}^{\infty} \frac{(\zeta(2) - \zeta_n(2))^2}{n} = 5\zeta(2)\,\zeta(3) - 9\zeta(5),$$

$$\sum_{n=1}^{\infty} \left(\bar{\zeta}(2) - \bar{\zeta}_n(2)\right)^2 = 3\zeta(2)\ln 2 - \frac{9}{4}\zeta(3) - \frac{5}{8}\zeta(4),$$

$$\sum_{n=1}^{\infty} \frac{(\zeta(2) - \zeta_n(2))\,(\zeta(3) - \zeta_n(3))}{n} = \zeta^2(3) - \frac{61}{48}\zeta(6),$$

$$\sum_{n=1}^{\infty} (\zeta(2) - \zeta_n(2))^3 = 9\zeta(2)\,\zeta(3) - \frac{35}{8}\zeta(6) - \frac{25}{2}\zeta(5),$$

$$\sum_{n=1}^{\infty} (\zeta^\star(2,1) - \zeta_n^\star(2,1))^2 = \frac{15}{2}\zeta(5) + 3\zeta(2)\,\zeta(3) - 4\zeta^2(3),$$

$$\sum_{n=1}^{\infty} \frac{\zeta^2(3) - \zeta_n^2(3)}{n} = -\frac{73}{4}\zeta(7) + \frac{3}{2}\zeta(3)\,\zeta(4) + 10\zeta(2)\,\zeta(5),$$

$$\sum_{n=1}^{\infty} (\zeta(2) - \zeta_n(2))\left(\bar{\zeta}(2) - \bar{\zeta}_n(2)\right) = \frac{3}{2}\zeta(2)\ln 2 - \frac{5}{4}\zeta(4) - \frac{3}{8}\zeta(3),$$

$$\sum_{n=1}^{\infty} (\zeta(2) - \zeta_n(2))\,(\zeta(3) - \zeta_n(3))\,(-1)^{n-1} = \frac{9}{16}\zeta(2)\,\zeta(3) - \frac{31}{32}\zeta(5),$$

$$\sum_{n=1}^{\infty} (\zeta(2) - \zeta_n(2))^2\,(\zeta(3) - \zeta_n(3)) = \frac{7}{6}\zeta(6) + \frac{3}{2}\zeta^2(3) - \frac{5}{2}\zeta(3)\,\zeta(4),$$

$$\sum_{n=1}^{\infty} (\zeta^\star(2,1) - \zeta_n^\star(2,1))\,(\zeta^\star(3,1) - \zeta_n^\star(3,1)) = -\frac{1}{3}\zeta(6) + 3\zeta^2(3) - \frac{5}{2}\zeta(3)\,\zeta(4),$$

$$\sum_{n=1}^{\infty} (\zeta(2) - \zeta_n(2))^2(-1)^{n-1} = 4\mathrm{Li}_4\left(\frac{1}{2}\right) - \frac{29}{8}\zeta(4) - \zeta(2)\ln^2 2 + \frac{1}{6}\ln^4 2 + \frac{7}{2}\zeta(3)\ln 2,$$

$$\sum_{n=1}^{\infty} \frac{\zeta_n(2)\,\zeta^2(3) - \zeta(2)\,\zeta_n^2(3)}{n} = -\frac{73}{4}\zeta(2)\,\zeta(7) + \frac{21}{8}\zeta(3)\,\zeta(6) + 25\zeta(4)\,\zeta(5) - \zeta^3(3),$$

$$\sum_{n=1}^{\infty}\left(\bar{\zeta}\left(2\right)-\bar{\zeta}_{n}\left(2\right)\right)\left(\bar{\zeta}\left(3\right)-\bar{\zeta}_{n}\left(3\right)\right)=\frac{21}{16}\zeta\left(4\right)+\frac{1}{2}\zeta\left(2\right)\ln^{2}2-\frac{1}{12}\ln^{4}2-2\mathrm{Li}_{4}\left(\frac{1}{2}\right)-\frac{3}{8}\zeta\left(2\right)\zeta\left(3\right).$$

Acknowledgments. The authors would like to thank the anonymous referee for his/her helpful comments, which improve the presentation of the paper.

References

[1] George E. Andrews, Richard Askey, Ranjan Roy. *Special Functions*. Cambridge University Press., 2000: 481-532.

[2] David H. Bailey, Jonathan M. Borwein and Roland Girgensohn. *Experimental evaluation of Euler sums*. Experimental Mathematics., 1994, **3**(1): 17-30.

[3] David Borwein, Jonathan M. Borwein and Roland Girgensohn. *Explicit evaluation of Euler sums*. Proc. Edinburgh Math., 1995, **38**: 277-294.

[4] Jonathan M. Borwein, David M. Bradley, David J. Broadhurst, Petr. Lisonk. *Special values of multiple polylogarithms*. Trans. Amer. Math. Soc., 2001, **353**(3): 907-941.

[5] J.M. Borwein, R. Girgensohn. *Evaluation of triple Euler sums*, Electron. J. Combin., 1996: 2-7.

[6] Minking Eie, Chuan-Sheng Wei. *Evaluations of some quadruple Euler sums of even weight*. Functions et Approximatio., 2012, **46**(1): 63-67.

[7] Philippe Flajolet, Bruno Salvy. *Euler sums and contour integral representations*. Experimental Mathematics., 1998, **7**(1): 15–35.

[8] O. Furdui. *Limits Series and Fractional Part Integrals: Problems in Mathematical Analysis, Problem Books in Mathematics*. Springer, London., 2013: 171-240.

[9] O. Furdui, A. Sîntămărian. *Quadratic series involving the tail of* $\zeta(k)$. Integral Transforms Spec. Func., 2015, **26**: 1-8.

[10] O. Furdui, C. Valean. *Evaluation of series involving the product of the tail of* $\zeta(k)$ *and* $\zeta(k+1)$. Mediterr. J. Math., 2016, **13**: 517-526.

[11] M. E. Hoffman. *Multiple harmonic series*. Pacific J. Math., 1992, **152**: 275-290.

[12] M. E. Hoffman. *Sums of Products of Riemann Zeta Tails*. Mediterr. J. Math., 2016: 1-11.

[13] I. Mezö. *Nonlinear Euler sums*. Pacific J. Math., 2014, **272**: 201-226.

[14] A. Sofo. *Quadratic alternating harmonic number sums*. J. Number Theory., 2015, **154**: 144-159.

[15] A. Sofo, H.M. Srivastava. *Identities for the harmonic numbers and binomial coefficients*. Ramanujan J., 2011, **25**: 93-113.

[16] A. Sofo. *Harmonic sums and integral representations*. J. Appl. Anal., 2010, **16**: 265-277.

[17] Ce Xu, Yuhuan Yan, Zhijuan Shi. *Euler sums and integrals of polylogarithm functions*. J. Number Theory., 2016, **165**: 84-108.

[18] Ce Xu, Mingyu Zhang, Weixia Zhu. *Some evaluation of q-analogues of Euler sums*. Monatshefte Für Mathematik ., 2016: 1-19.

[19] D. Zagier.*Values of zeta functions and their applications*. First European Congress of Mathematics, Volume II, Birkhauser, Boston., 1994, (120): 497-512.

[20] D. Zagier. *Evaluation of the multiple zeta values* $\zeta(2,...,2,3,2,...,2)$. Annals of Mathematics., 2012, **2**(2): 977-1000.